



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).


# MISCELLANEOUS.

58. Proposed by EDMUND FISH, Hillsboro, Ill.

The longest noonday winter shadow of an upright object is found to be seven times as long as the shortest summer shadow of the same object. Required the latitude of the place.

I. Solution by S. HART WRIGHT, A. M., M. D., Ph. D., Penn Yan, N. Y.

In the right plane triangles  $ABC$  and  $APC$ , let the vertical  $AC$  (=unity) be the rod that casts a shadow from  $C$  to  $B$ , and from  $C$  to  $P$ , when the sun is at  $S$  and  $D$ . Extend  $CA$  to the zenith  $Z$ ,  $BA$  to  $S$  and  $PA$  to  $D$ . Bisect  $DAS$  with  $QA$ . Let  $\angle BAC = \chi = \angle ZAS$ .  $DAS$  is double the obliquity of the ecliptic  $= 2\delta = \angle PAB = 46^\circ 54' 30'' = v$ .  $DAQ = SAQ = \delta$ , and  $Q$  must be on the equator, and  $QAZ$  = the required latitude  $= \lambda$ .  $CP = 7CB$ . Put  $7 = m$ , and  $\tan^{-1}m = \beta = 81^\circ 52' 12''$ . We have  $CB = \tan \chi$ , and  $CP = \tan(\chi + v)$ , and  $m \tan \chi = \tan(\chi + v)$ , a trigonometric equation, from which we derive  $\sin(2\chi \pm v) = \cot(\beta \mp 45^\circ) \sin v$ . Four values of  $\chi$  result, the upper signs giving the only acceptable value of  $\chi = 14^\circ 57' 30''$ . The other signs make the  $\angle PAC > 90^\circ$ . Now  $\lambda = \chi + \delta = 38^\circ 24' 45''$  north or south, as the seasons are interchangeable on each side of the equator.



II. Solution by G. B. M. ZERR, A. M., Ph. D., President and Professor of Mathematics, The Russell College, Lebanon, Va.

Let  $AC$  be the upright object length  $= l$ ,  $CP$  the earth,  $SB$  the summer sun,  $DP$  the winter sun. Let  $\phi$  = latitude,  $\delta$  = north,  $-\delta$  = south declination of the sun, and let the winter shadow be  $n$  times as long as the summer shadow.

Then  $\angle CAB = (\phi - \delta)$ ,  $\angle CAP = (\phi + \delta)$ ,  $CP = n \cdot CB$ ,  $CP = l \tan(\phi + \delta)$ ,  $CB = l \tan(\phi - \delta)$ .

$$\therefore l \tan(\phi + \delta) = n l \tan(\phi - \delta).$$

$$\therefore (n + 1) \tan^2 \phi \tan \delta - (n - 1) \tan \phi \sec^2 \delta + (n + 1) \tan \delta = 0.$$

$$\tan^2 \phi - \frac{2(n - 1) \tan \phi}{(n + 1) \sin 2\delta} + 1 = 0.$$

$$\therefore \tan \phi = \frac{(n - 2) \pm \sqrt{(n - 1)^2 - (n + 1)^2 \sin^2 2\delta}}{(n + 1) \sin 2\delta} \dots \dots \dots (A).$$

Now  $\delta = 23^\circ 27' 30''$ ,  $n = 7$ .  $\therefore \tan \phi = .793428$  or  $1.26035$ .

$\therefore \phi = 38^\circ 25' 46''$  or  $51^\circ 34' 14''$ .

The two values of  $\tan \phi$  are equal when  $n = (1 + \sin 2\delta) / (1 - \sin 2\delta)$ .

$\therefore n = 6.4118$ ,  $\phi = 45^\circ$ . When  $\phi = \delta$ , and  $\phi + \delta = 90^\circ$ ,  $n$  is infinite.

In the first case the summer shadow is zero and the winter shadow is fi-

nite ; in the second case, the winter shadow is infinite and the summer shadow is finite.

In formula (A),  $\delta$  and  $n$  can have any values within proper limits.

Also solved by W. W. LANDIS, and J. SCHEFFER.

59. Proposed by J. A. CALDERHEAD, M. Sc., Professor of Mathematics, Curry University, Pittsburg, Pa.

When a cylindrical china jar, standing upon the ground, receives the sun's rays obliquely, a bright curve is observed to form itself at the bottom of the jar, and it is found that the shape and dimensions of this curve are not affected by the varying elevations of the sun: account for this latter circumstance, and determine the nature of the bright curve. [From *Parkinson's Optics*.]

Solution by ALFRED HUME, C. E., D. Sc., Professor of Mathematics, University of Mississippi, University, Mississippi.

All rays striking any element of the cylindrical surface lie in a vertical plane. Their reflections form the other face of the dihedral angle whose bisector passes through the axis of the cylinder. These reflected rays intersect the base of the cylinder in a straight line. There is thus formed a system of lines, and the bright curve observed is their envelope. The altitude of the sun does not affect the position of the vertical planes; and, therefore, the intersections with the bottom of the jar are unchanged, and the continual intersection of the consecutive lines so formed produces a curve invariable as to its shape and size.

The bright curve is the caustic by reflection for the circle, the incident rays being parallel. The following general property of caustics by reflection for parallel rays is established in *Price's Infinitesimal Calculus*: "The distance from the incident point in the reflecting curve to the point of intersection of two consecutive reflected rays, is equal to one-fourth of the chord of the circle of curvature at the point of incidence which is parallel to the incident ray."

A. Take the center of the circle as the origin, the  $X$ -axis parallel to the incident rays, the  $Y$ -axis perpendicular to them.

Let  $AB$  be an incident ray,  $BC$  its reflection, the angle between them being  $2\theta$ . Take  $BP$  along  $BC$  equal to one-half of  $DB$ ,  $D$  being the intersection of  $AB$  with the  $Y$ -axis. Then, according to the principle quoted above,  $P$  is a point of the caustic. To find the locus of  $P$ : Draw  $OB$ , denoting it by  $a$ . From  $P$  drop a perpendicular to  $AB$  meeting it at  $H$ . Denoting the coördinates of  $P$  by  $x$  and  $y$ ,

$$x = DB - HB = a \cos \theta - \frac{1}{2} a \cos \theta \cos 2\theta,$$

$$y = OD - PH = a \sin \theta - \frac{1}{2} a \cos \theta \sin 2\theta.$$

From these  $x = \frac{1}{2} a \cos \theta - \frac{1}{2} a \cos 3\theta$ ,

$$\text{and } y = \frac{1}{2} a \sin \theta - \frac{1}{2} a \sin 3\theta.$$

These may be written

